

Agarikon.1 and Agarikon Plus Affect Cell Cycle and Induce Apoptosis in Human Tumor Cell Lines

Jakopovich Boris, Jakopovich Ivan,
Jakopovich Neven

Dr Myko San – Health from
Mushrooms

INDEX

- Introduction
- Materials and Methods
- Results
- Conclusions

INTRODUCTION

- Continuation of the study on 6 blended mushroom products and 3 single extracts on 4 human tumor cell lines (Durgo, Jakopovich 2013)
- Neutral Red and MTT proliferation assays demonstrate that blended extracts cause increased tumor cell membrane and mitochondria damage
- This study concentrates on the mechanisms; effects on the cell cycle and inducing apoptosis

Medicinal mushroom metabolites can interfere and modulate all processes related to the 8 steps of carcinogenesis (Petrova 2012):

- inflammation
- cancer cell proliferation
- adhesion
- apoptosis
- angiogenesis
- gene expression
- invasiveness
- metastasis

Apoptosis - programmed cell death - is a crucial tumor suppression mechanism

- eliminates cells with extensive DNA damage (potentially leading to cancer)
- differentiation - cell growth – apoptosis balance

Purpose of the Study

Investigate:

A. proliferation (cell cycle disturbance)

B. induction of apoptosis

- medicinal mushroom extract blends Agarikon Plus and Agarikon.1

- concentration response

- 24 and 48 hour response

- 2 human tumor cell lines: H460 (lung carcinoma) and Caco-2 (colon carcinoma)

- Camptothecin, referent antitumor compound; cytotoxic dose 10 μM used

MATERIALS AND METHODS

Cell lines

- H460 (large cell lung carcinoma)
- Caco-2 (colorectal adenocarcinoma)

Tested extract blends

- Agarikon.1 tablets
- Agarikon Plus

Experimental methods

- Proliferation Assay by MTT
- Cell Cycle Analysis by flow cytometry
- Annexin V Assay for Apoptosis Induction Detection
- Western Blot Analysis

TESTED PRODUCTS

- Agarikon.1 tablets (AG.1)
- proprietary mushroom extract blend from Dr Myko San company
- *Ganoderma lucidum*, *Lentinus edodes*, *Grifola frondosa*, *Pleurotus ostreatus*, *Agaricus brasiliensis*
- registered med. mushroom supplement
- recommended treatment dose: ~0.1 g/kg bodyweight per day of soluble polysaccharides



- Agarikon Plus extract blend (AG+)
- proprietary mushroom extract blend from DMS
- 10 medicinal mushroom species (inc. *G. lucidum*, *L. edodes*, *G. frondosa*, *P. ostreatus*, *A. brasiliensis*)
- in liquid form
- Recommended treatment dose: $\sim 0.16\text{g/kg}$ BW per day of soluble polysaccharides



1 Proliferation Assay

- Cells cultured as monolayers, plated in parallel on day 0, at 3×10^3 cells/well (H460) and 7×10^3 cells/well (Caco-2), depending on doubling times
- AG.1 and AG+ added at 0.001, 0.01, 0.1, 1 and 10 mg/ml concentrations (stock solution for both 40 mg/ml, and 4×10^{-3} M/DMSO for camptothecin)
- We used MTT assay to evaluate cell growth rate after 72 hours (absorbance was measured at 570 nm)

2 Cell Cycle Analysis

- seeded at 1×10^5 cells/well (H460) and 2×10^5 cells/well (Caco-2), depending on the doubling times
- After 24 hours, AG.1 and AG+ applied at concentrations 0.1 mg/ml and 1 mg/ml; camptothecin ($10 \mu\text{M}$) for positive control
- After the incubation period, cells were trypsinized, washed with Phosphate Buffer Saline (PBS); stained with propidium iodide (PI) and analyzed on FACScalibur flow cytometer
- Ratio of cells in each cell cycle phase was determined by analyzing the DNA histograms using ModFit LTTM software

3 Annexin V Assay for Apoptosis Induction Detection

- same concentrations used; 0.1 and 10 mg/ml
- cell populations were gated into regions corresponding to live, early apoptotic and late apoptotic/necrotic cells

Annexin V	PI	Cell Region
-	-	Live cells
+	-	Early apoptotic
+	+	Late apoptotic /necrotic

4 Western Blot Analysis

- mushroom extracts (0.1 and 1 mg/ml) were added to well plates after 24 hours
- total proteins were measured using BCA Protein Assay Reagent, separated by SDS-polyacrylamid gel electrophoresis and transferred to PVDF membrane → probing with anticaspase 3, anti-p53, and anti-p21 primary antibodies
- equal loading confirmed using anti-tubulin primary antibody

RESULTS


Proliferation Assay

- Agarikon Plus (strong effect at 10 mg/ml, GI₅₀ ≈ 2-3 mg/ml) and Agarikon.1 inhibit the growth of both tumor cell lines
- H460 cells are more resistant to Agarikon.1 (approaching GI₅₀ above 10 mg/ml mass concentration)


GI ₅₀ ^a (mg/ml)		
Test agent	Cell lines	
	Caco-2	NCI-H460
Agarikon Plus	1.9 ± 0.1	3.4 ± 1
Agarikon.1	1.6 ± 0.3	≥10

^a GI₅₀; growth inhibition 50 - the concentration that causes 50% growth inhibition


- Concentration-response curves showing growth inhibition of H460 and Caco-2 cell in vitro after 72 hours after adding **Agarikon.1**, **Agarikon Plus**, and **camptothecin**.



Agarikon Plus



Camptothecin




- expected Camptothecin curve confirms valid measurements (positive control)

Cell Cycle Analysis


- Distribution of **H460** cells by cell cycle phase: G0/G1, S, G2/M; and sub G1 (dead/apoptotic cells)
- treated with Agarikon Plus, Agarikon.1 (at 0.1 and 10 mg/ml), camptothecin at 10 μ M.
- measured by flow cytometry at 24 and 48 hours

H460 - 24h




- no significant influence on cell cycle
- moderate increase in apoptotic/dead cells for AG+ at higher concentrations

H460 - 48h




- AG+ and AG.1 (1mg/ml) induce accumulation of cells in G1, reduction in S, increase in sub G1 (apoptotic/necrotic)

Caco-2 - 24h



- AG+ (1mg/ml): reduced G1, increased S
- no significant sub G1 influence

Caco-2 - 48h




- AG+, AG.1 (1 mg/ml) reduced G1, increased S phase
- no significant sub G1 influence

Apoptosis Induction Detection by Annexin V Assay


- Ratio of H460 cells in early or late apoptosis, obtained by co-staining with FITC-labeled annexin V and propidium iodide (PI) and analyzed by flow cytometry.

H460 - 24h




- AG+ (1 mg/ml): moderate increase in early apoptotic cells

H460 - 48 h




- AG+ (1 mg/ml): larger increase in late apoptotic/necrotic cells

Caco-2 - 24h



- no significant influence, AG+ (1mg/ml)
moderate early apoptotic cell increase


Caco-2 - 48h



- AG+ (1 mg/ml): moderate increase in induced early apoptosis


Apoptosis induction detection by caspase-3 cleavage assessment

- Appearance of the 17-kDa subunit (caspase-3 p17) – a major cleaved product of the 32-kDa zymogen procaspase-3 - confirms caspase-3 activation which marks the induction of apoptosis.




The effect of AG+ and AG.1 on cleavage of procaspase-3 in H460 cells (24; 48 hours)

- A. AG+ at 0.1 mg/ml (lanes 1;6), 1 mg/ml (lanes 2;7).
- B. AG.1 at 0.1 mg/ml (lanes 3;8), 1 mg/ml (lanes 4;9)
- C. **Camptothecin** at 10 μ M (lanes 5;10)




The effect of AG+ and AG.1 on cleavage of procaspase-3 in **Caco-2** cells (24; 48 hours)


- A. AG+ at 0.1 mg/ml (lanes 1;6), 1 mg/ml (lanes 2;7).
- B. AG.1 at 0.1 mg/ml (lanes 3;8), 1 mg/ml (lanes 4;9)
- C. **Camptothecin** at 10 μ M (lanes 5;10)



Influence of AG+ and AG.1 on p53 and p21 protein expression

- Agarikon.1, in both concentrations, induced mild p53 protein expression in H460 after 24 hours (A; lanes 3 and 4)





- Both Agarikon Plus (lane 2) and Agarikon.1 (lane 4) at the concentration of 1 mg/ml induced a moderate increase in the expression of both p53 and p21 after 24 hours in H460

CONCLUSIONS

- Agarikon Plus and Agarikon.1 possess antiproliferative, mainly cytostatic activity, on H460 and Caco-2 cells, in the concentration range 1-10 mg/ml
- Both induce cell cycle perturbations, by delaying the progress through the G1 and S phase
- This points to disturbances occurring before or during DNA replication (confirmed by increase in both p53 and p21 protein expression)

CONCLUSIONS (II)

- Although a modest induction of early (after 24 hours) and late (48 hours) apoptosis was noticed by annexin V test, no processing (cleavage) of caspase-3 was detected
- More-pronounced antiproliferative activity (MTT) of tested agents towards Caco-2 line at maximal concentration (10 mg/ml) points to a non-specific cytotoxic effect

Acknowledgements

Dr. Marijeta Kralj

Dr. Ana-Matea Mikecin

Dr. Damir Kralj

Biozyne, Bijenicka c. 54, 10000 Zagreb,
Rudjer Boskovic Institute

Thank you for your attention!

- Send correspondence to:
borisjakopovic@yahoo.com

Percentage of Growth definition

The percentage of growth (PG) expressions:

1. For $(A_t - A_0) \geq 0$:

$$PG = 100 * (A_t - A_0) / (A_c - A_0)$$

2. For $(A_t - A_0) < 0$:

$$PG = 100 * (A_t - A_0) / A_0$$

A_0 ... avg. absorbance before exposure

A_t ... avg. absorbance after test (72 h)

A_c ... avg. absorbance after 72 h, no exposure to test compound

Camptothecin Proliferation Assay

